arXiv:1301.6954v1 [physics.optics] 29 Jan 2013

Twisting Electromagnetic Fields with Singular Transformation Optics

Yong-Liang Zhang[1], Li-Na Shi[2], Xian-Zi Dong[1], Zhen-Sheng Zhao[1] and Xuan-Ming Duan[1]*
[1] *Key Laboratory of Functional Crystal and Laser Technology,*
Technical Institute of Physics and Chemistry, Chinese Academy of Science, Beijing 100190, China and
[2] *Key Laboratory of Nano-Fabrication and Novel Devices Integrated Technology,*
Institute of Microelectronics, Chinese Academy of Sciences, Beijing 100029, China

Transformation optics is a powerful method for manipulating the electromagnetic fields with metamaterials which resembles a optical Riemannian space for photons. Analogous to the theory of topological defects, we propose a geometric extension of transformation optics by introduction of linear topological defect which produces spatial torsion. In order to illustrate the optical effect of singular transformation optics, we present an example of application to control the twisting degree of freedom of light, the orbital and spin angular momentum of light with an optical cosmic string. Our results provide a systematic and versatile framework which may open intriguing venues in angular momentum engineering of light with artificial metamaterials.

Introduction.—In recent years, transformation optics (TO) has attracted a great deal of attention because it promises a variety of intriguing applications from invisibility cloaking, electromagnetic wormholes to broadband light harvesting [1–5, 8–12]. The underlying physical idea of TO stems from the formal invariance of the governing Maxwell's equations [1] and Helmholtz equation [2] under a spatial coordinate transformation. The constitute parameters of materials ε and μ are renormalized [6] by the transformation's Jacobian matrix which represents the metric tensor of the transformed space. Alternative approaches have also been proposed, based on, e.g. space-time transformations [13], direct fields transformations [14], and spectra domain [15] transformations which generalize the transformation media to include moving, bianisotropic and nonreciprocal media.

So far the rapid growing development has proved TO as a powerful and well-established tool for the systematic design of artificial metamaterials with specified electromagnetic energy flow distributions, and the same idea has made important contributions to advancing the manipulation of sound wave [16], matter wave [17], dc electric current [18] and even heat flux [19]. It is usually accepted that the inhomogeneous optical material acts as a pseudo-Riemannian manifold, where the propagation of photons follows null geodesic determined by local spatial curvature [7, 20–22]. However, the vector nature of electromagnetic waves require some extensions of the standard TO to include the twisting degree of freedom of light, such as optical spin and orbital angular momentum (AM). Horsley [30] has earlier shown that an inhomogeneous isotropic Tellegen medium can be interpreted as a Weitzenböck space where the chiral parameter is assigned with the contorsion tensor.

Although the approaches in [14, 30] seem to capable to include chiral metamaterial, no further clues with geometric intuitive were given to extend the standard TO to embody the twisting degree of freedom of light. Moreover, the vast majority of reported TO designs are based holonomic coordinate transformation which preserve the topology between the virtual and physical spaces. Here we explore the extension of TO of Pendry and Leonhardt to include spatial torsion and investigate the physical effects of torsional media on the propagation of electromagnetic waves, focusing on the generation and physical effects of electromagnetic torsion by anholonomic coordinate transformation. Inspired by the theory of topological defects in crystalline solid or gravitation [24–29], we introduced torsion in terms of the spatial gradient of the inverse lattice deformation in subwavelength scale and naturally arises in Maxwell equations in transformed space as magnetoelectric coupling terms. In addition, anholonomic coordinate transformation in macroscopic scale can transform light beams with planar wave front to beams with helical wave front. Numerical fullwave calculations based on a specific example of linear screw dislocations confirm the theoretical predictions. Our model not only extends the geometrical and topological aspects of TO but also provides an experimental approach for simulating of classical and quantum physics in curved torsional space in optical laboratory.

Topology and geometry of TO.—Similar to the elastic deformation of perfect crystal, method of TO relies on a spatial coordinate transformation from a prescribed domain in Cartesian coordinate system to a deformed one. The topology of TO will change if the spatial continuity is broken under the deformation. Our proposed approach is based on the anholonomic coordinate transformation in the spatial domain, which is often indicated the emergence of topological defects.

As a starting point, we considering a coordinate transformation from an initial region $\mathcal{D}$ in the flat space, identified by $\mathbf{r} \equiv \{x^a|(x, y, z)\}$ to a transformed region $\mathcal{D}'$ in physical space, identified by prime coordinates $\mathbf{r}' \equiv \{x^\mu|(x', y', z')\}$. Here the Latin and Roman letter are used for the superscript to coincide with the symbol of tetrad theory of gravity. The action of coordinate transformation on point $\mathbf{r}$ is generally described by a locally special Euclidean group $SE(3)$ [31], i.e.

---

* Corresponding author: xmduan@mail.ipc.ac.cn

$$\mathbf{r}' = \boldsymbol{O}(\mathbf{r})\mathbf{r} + \boldsymbol{T}(\mathbf{r}) \tag{1}$$

where $\boldsymbol{O} \in SO(3)$ and $\boldsymbol{T} \in T(3)$. Therefore, there are six degrees of freedom for coordinate transformation describing rotational and translational symmetries in $\mathbb{R}^3$. From the view point of topology, the virtual space and the transformed space are generally assumed diffeomorphic which means the two space are topologically equivalent. The material interpretation of TO regard the domains $\mathcal{D}$ and $\mathcal{D}'$ as some elastic objects, which carry systems of coordinates together with electric displacement field lines in an absolute Euclidean space, the transformed region $\mathcal{D}'$ is just a deformed body with curved coordinated lines which is labeled by a metric tensor $g_{\mu\nu}(\mathbf{r}) = \delta_{ab}\frac{\partial x^a}{\partial x^\mu}\frac{\partial x^b}{\partial x^\nu}$, and the constitutive parameters of transformed media is given by

$$\chi^{\mu\nu} = \sqrt{\det(g)}g^{\mu\nu}; \quad \mathbf{G}'_\mu = e^a_\mu \mathbf{G}_a, \tag{2}$$

with $\det(\cdot)$ denoting the matrix determinant and χ, $\mathbf{G}$ representing (ε, μ) and ($\mathbf{E}$, $\mathbf{H}$), respectively. Here and below, we use the Einstein summation convention in which repeated indices are summed.



FIG. 1. Schematic illustration of anholomonic coordinate transformation in a discrete lattice model. A closed loop *ABCDA* in is mapped to a open curve *A1B1C1D1*. The failure of the closed loop indicates a line defect along z-axis.

The key point for going beyond Riemannian geometry comes from the observation that the Jacobian matrix constructed the tetrad fields in the affine space, i.e. $\mathbf{e}_{(\mu)} = \partial\mathbf{r}/\partial x^\mu$, which related the basis vector $\mathbf{e}_{(\mu)}$ in $\mathcal{D}'$ and the basic vector $\mathbf{e}_{(a)}$ in $\mathcal{D}$,

$$\mathbf{e}_{(\mu)} = e^a_\mu \mathbf{e}_{(a)} \tag{3}$$

where $\mu, a = 1, 2, 3$ are the spatial indices. Torsion tensor is defined in terms of the asymmetry part of the affine connection $T^\lambda_{\mu\nu} = (\Gamma^\lambda_{\mu\nu} - \Gamma^\lambda_{\nu\mu})/2$, where the affine connection is given by $\Gamma^\lambda_{\mu\nu} = \mathbf{e}^\lambda \cdot \partial_\mu \mathbf{e}_\nu$. Torsion vanishes for trivial coordinate transformation which create effective Riemannian medium for the photons. By definition, a transformed space with torsion demands that the second partial derivatives of the coordinate transformation don't commutable [29], that implies the coordinate transformation is subject to an anholonimic constraint which breaks the translational symmetry,

$$\frac{\partial^2 x^a}{\partial x^\mu \partial x^\nu} - \frac{\partial^2 x^a}{\partial x^\nu \partial x^\mu} \neq 0 \tag{4}$$

Inequality (4) means the violation of Schwartz integrable condition [29]. From a geometric point of view, the constrained coordinate transformation leads to a topology transition form the auxiliary space $\mathcal{D}$ to $\mathcal{D}'$. In other words, the transformed space $\mathcal{D}'$ must contained some topological defects, such as dislocations in continuum limit.

Unlike the well known meaning of curvature which characters how a tangent vector is rotated along a closed loop, torsion tensor measures the degree of how the tangent spaces is *twisted* about a curve when they are parallel transported,

$$d\mathbf{E}'^\lambda = (R^\lambda_{\rho\mu\nu}\mathbf{E}'^\rho + T^\lambda_{\mu\nu})dS^{\mu\nu} \tag{5}$$

If we define the displacement vector from a perfect lattice to a deformed lattice by $\mathbf{u} = \mathbf{r} - \mathbf{r}'$. Encircling the defect with an arbitrary closed contour C, the discontinuity of $\mathbf{u}$ is given by

$$\oint_C (dx \cdot \nabla)\mathbf{u} = \mathbf{b} + \boldsymbol{\Omega} \times (\mathbf{r} - \mathbf{r_0}) \tag{6}$$

where the Frank vector $\boldsymbol{\Omega}$ represents a rotational action, the Burgers vector $\mathbf{b}$ denotes the translational action and $\mathbf{r_0}$ the reference point. Figure 1 schematic shows the failure of parallelogram around a screw dislocation. The emergence of topological defects allows us to distinguish different loops connecting arbitrary two points in transformed space via homotopy classes. For the case of transformed media with a single linear topological defect, the fundamental group is $\pi_1(S^1) = \mathbb{Z}$.

Physical effects of torsion.—Physically, the permittivity ε and permeability μ, i. e. the metric tensor $\mathbf{g}$, fixes the causal structure of the space-time as well as the local optical length; the connections describes the parallel transport of free photons. The idea to relate vortices with optical dislocations seems intuitively obvious since vortices are also known as optical phase dislocations with an $\exp(il\theta)$ azimuthal phase dependence. From TO, the electric field $\mathbf{E}'$ in transformed medium is related the original $\mathbf{E}$ via

$$\mathbf{E}'_\mu = e^a_\mu \mathbf{E}_a \tag{7}$$

Consider a plane wave in virtual space is transformed under the action of e^a_μ, the contour integral of $\mathbf{E}'$ around z axis over a closed loop, based on Stokes theorem, can be expressed as

$$\begin{aligned}\oint_C dx^\mu \mathbf{E}'_\mu &= (\int_S ds^{\mu\nu} T^a_{\mu\nu})\mathbf{E}_a \\ &= \mathbf{b}^a E_a\end{aligned} \tag{8}$$

(8) indicate that a macroscale topological defect with torsion can convert a plane wavefront to a dislocated one. Specifically, a screw dislocation will leading to a optical vortex with topological invariant **b**.

To see how torsion is linked with magnetoelectric coupling in constitutive relations, we follows the idea of standard TO based on formal invariance. Starting from Maxwell equations in vacuum, anholonomic coordinate transformation leading to non-vanishing terms $\epsilon_{\mu\nu\lambda}(\partial^2 x_a/\partial x_\nu \partial x_\lambda)E_a$. A simple calculations give out the following constitutive relations,

$$\begin{aligned} \mathbf{D} &= \varepsilon\mathbf{E} + i\kappa\mathbf{H} \\ \mathbf{B} &= \mu\mathbf{H} - i\kappa\mathbf{E} \end{aligned} \tag{9}$$

with $\varepsilon = \mu = \sqrt{det(g)}g^{\mu\nu}$ and $\kappa^{\mu\rho} = \frac{1}{2\omega}\epsilon^{\mu\nu\lambda}T^{\rho}_{\nu\lambda}$.

Consider the local density of optical chirality C which was firstly introduced by Lipkin [32, 33]

$$C = \frac{\varepsilon_0}{2}\mathbf{E}'\cdot\nabla\times\mathbf{E}' + \frac{1}{2\mu_0}\mathbf{B}'\cdot\nabla\times\mathbf{B}' \tag{10}$$

According to (2), we have

$$C = \varepsilon_{\mu\nu\lambda}g_{\mu\rho}T^{\rho}_{\nu\lambda}U \tag{11}$$

where $U = \frac{1}{2}(\varepsilon_0\mathbf{E}^2 + \frac{1}{\mu_0}\mathbf{H}^2)$ is the energy density of light field.

As an illustrative example, we consider a curved space with a topological defect which is described by the following transformation in cylindrical coordinate[27]

$$r' = r; \quad \theta' = \alpha\theta; \quad z' = z + \frac{\beta}{\alpha}\theta, \tag{12}$$

where $\alpha = 1 - 4\mu$, $\beta = b/2\pi$ is the Burgers vector. Obviously, this line element turns to be vacuum for $\alpha = 1$ and $\beta = 0$, a conical space for $\beta = 0$, and a screw dislocation or a static chiral cosmic string for $\alpha = 1$.

Through this transformation, we can obtain the following constitutive parameters for the material

$$\varepsilon = \mu = \frac{M}{\alpha^3} \tag{13}$$

with

$$M = \begin{pmatrix} sin^2\theta + \alpha^2 cos^2\theta & sin\theta cos\theta[\alpha^2 - 1] & \gamma sin\theta \\ & cos^2\theta + \alpha^2 sin^2 & -\gamma cos\theta \\ & & \alpha^2 + \gamma^2 \end{pmatrix}$$

The material parameters (13) describe an optical medium with the material distribution breaking translational and/or rotational symmetries. To explore the optical properties, we perform fullwave numerical simulations with finite element method (FEM). Figure 2 depicts the electromagnetic field distribution of a Gaussian beam incident on a cylinder of defect materials with the defect line is aligned with $z-$axis. It is shown the plane wavefront of a gaussian wave is titled in the defect media region, and the EM wave intensity in the defect core. As a result, a gaussian beam is converted to be an optical vortex which carrying orbital angular momentum.



FIG. 2. Propagation of a Gaussian beam along an optical line defect with (a) $\beta = 0.15, \alpha = 1$, (b) $\beta = -0.1, \alpha = 1$ and (c) $\beta = -0.1$, $\alpha = 1.3$.

For a specific parameter $\beta = 0$, coordinate transformation (1) represents a static cosmic string. In Fig. 3, we plot the scattering pattern of a vertically incident gaussian beam on a cylinder consist of 2D conical media, i. e. $\mu \neq 1$, and $\beta = 0$ with μ is the deficit angle. For $\mu = 0$, the conical medium account for air. When μ decreased, the incident beam is split to two beams with the split angle increased with the deficit angle. We noted the observed phenomena is an optical analogy of gravitation lensing effect by 'cosmic strings'.



FIG. 3. Scattering of Gaussian beams by cylinders with 2D conical geometry with different μ: (a,b) $\mu = -1.6$, (c,d) $\mu = -0.8$, (e,f) $\mu = -0.2$, (g,h) $\mu = 0$, (i,j) $\mu = 0.04$.

Conclusion.— We show that the geometry framework of transformation optics, the powerful methodology for engineering the electromagnetic fields with metamaterials, can be extended to Riemann-Cartan geometry. Relying on the anholonomic coordinate transformation, one can create optical analogy of topological defect and introduce torsion tensor in transformation media. In order to illustrate the optical effect of singular transformation optics, we present an example of application to controlling the angular momentum, both orbital and spin, of light with an optical cosmic string. Our results provide a systematic and versatile framework which may open intriguing venues in angular momentum engineering of light with artificial metamaterials.

This work was financially supported by the National Basic Research Program (2010CB934102); Natural Science Foundation of China (NSFC, No. 60907019, No. 61077028, No. 50973126); The International Cooperation Program under Grant (No. 2008DFA02050, No. 2010DFA01180) of Most of China.

---

[1] J. B. Pendry, D. Schurig and D. R. Smith, *Controlling electromagnetic fields*, Science **312**, 1780 (2006).

[2] U. Leonhardt, *Optical conformal mapping*, Science **312**, 1777 (2006).

[3] H. Y. Chen, C. T. Chan and P. Shen, *Transformation optics and metamaterials*, Nat. Mat. **9**, 387 (1010).

[4] D. Schurig, J. J. Mock, B. J. Justice, S. A. Cummer, J. B. Pendry, A. F. Starr and D. R. Smith, *Metamaterial electromagnetic cloak at microwave frequencies*, Science **314**, 977-980 (2006).

[5] Y. Lai, J. Ng, H. Y. Chen, D. Z. Han, J. J. Xiao, Z. Q. Zhang and C. T. Chan, *Illusion optics: the optical transformation of an object into another object,* Phy. Rev. Lett. **102**, 253902 (2009).

[6] J. Plebanski, *Electromagnetic waves in gravational fields*, Phys. Rev. **118**, 1396 (1960).

[7] U. Leonhardt and T. G. Philbin, *General relativity in electrical engineering*, New J. Phys. **8**, 247 (2006).

[8] A. Greenleaf, Y. Kurylev, M. Lassas and G. Uhlmann, *Electromagnetic wormholes and virtual magnetic monopoles from metamaterials,* Phy. Rev. Lett. **99**, 183901 (2007).

[9] M. Li, R. X. Miao and Y. Pang, *More studies on metamaterials mimicking de Sitter space,* Opt. Express *18*, 9026-9033 (2010).

[10] Y. Luo, D. Y. Lei, S. Maier and J. B. Pendry, *Broadband light harvesting nanostructures robust to edge bluntness,* Phys. Rev. Lett. **108**, 023901 (2012).

[11] R.T. Thompson, and J. Frauendiener, *Dielectric analog space-times*, Phys. Rev. D **82**, 124021 (2010).

[12] T. G. Mackay and A. Lakhtakia, *Towards a metamaterial simulation of a spinning cosmic string*, Phys. Lett. A **374**, 2305-2308 (2010).

[13] M.W. McCall, A. Favaro, P. Kinsler and A. Boardman, *A spacetime cloak, or a history editor*, J. Opt. **12**, 024003 (2011).

[14] S.A. Tretyakov, I.S. Nefedov, and P. Alitalo, *Generlized field-transforming metamaterials*, New J. Phys. **10**, 115028 (2008).

[15] G. Castaldi, V. Galdi, A. Alu, and N. Engheta, *Nonlocal transformation optics*, Phys. Rev. Lett. **108**, 063902 (2012).

[16] G.W. Milton, M. Briane, and J.R. Willis, *On cloaking for elasticity and physical equations with a transformation invariant form*, New J. Phys. **8**, 248 (2006).

[17] S. Zhang, D. A. Genov, C. Sun, and X. Zhang, *Cloaking of matter waves*, Phys. Rev. Lett. **100**, 123002 (2008).

[18] F. Yang, Z.L. Mei, T.Y. Jin, and T.J. Cui, *dc electric invisibility cloak*, Phys. Rev. Lett. **109**, 053902 (2012).

[19] S. Guenneau, C. Amra, and D. Veynante, *Transformation thermodynamics: cloaking and concentrating heat flux*, Opt. Express **20**, 8027-8218 (2012).

[20] K. Liu, C. Song and M. L. Ge, *The geodesic form of light-ray trace in the inhomogeneous media*, Opt. Express **17**, 11753-11767 (2009).

[21] R. A. Crudo and J. G. O'Brien, *Metric approach to transformation optics,* Phys. Rev. A **80**, 033824 (2009).

[22] V.H. Schultheiss, S. Batz, A. Szameit, F. Dreisow, S. Nolte, A. Tunnermann, S. Longhi, and U. Peschel, *Optics in curved space*, Phys. Rev. Lett. **105**, 143901 (2010).

[23] F. W. Hehl, P. V. Heyde and G. D. Kerlick, *General relativity with spin and torsion: Foundations and prospects,* Rev. Mod. Phys. **48**,393-416 (1976).

[24] K. Kondo, *On the geometrical and physical foundations of the theory of yielding,* in: *Proc. 2nd Japan Nat. Congr. Applied Mechanics,* Tokyo, 41-47 (1952).

[25] A. Kadic and D. G. B. Edelen, *A gauge theory of dislocations and disclinations,* (Springer-Verlag, Berlin-Heidelberg, 1983).

[26] M. O. Katanaev and I. V. Volvich, *Thoery of defects in solids and tree-dimensional gravity*, Ann. Phys. (N.Y.), **216**, 1-28 (1992).

[27] A. Vilenkin and E. P. S. Shellard, *Cosmic strings and other topological defects,* (Cambridge University Press, 1994).

[28] D. V. Gal'tsov and P. S. Letelier, *Spinning strings and cosmic dislocations*, Phys. Rev. D **47**, 4273-4276 (1993).

[29] H. Kleinert, *Multivalued fields in condensed matter, electromagnetism, and Gravitation* (World Scientific, Singapore 2008).

[30] S. A. R. Horsley, *Transformation optics, isotropic chiral media and non-Riemannian geometry*, New J. Phys. **13**, 053053 (2011).

[31] R.A. Puntugam, and H.H. Soleng, *Volterra distortions, spinning strings, and cosmic defects*, Class. Quantum Grav. **14**, 1129-1149 (1997).

[32] D. Lipkin, *Exsitence of a new conservation law in electromagnetic theory,* J. Math. Phys. **5**, 696 (1964).

[33] Y.Q. Tang, and A.E. Cohen, *Optical chirality and its interaction with matter,* Phys. Rev. Lett. **104**, 163901 (2010).

[34] M. Schäferling, D. Dregely, M. Hentschel, and H. Giessen, *Taioring enhanced optical chiality: design principles for chiral plasmonic nanostructures,* Phys. Rev. X **2**, 031010 (2012).

[35] Y.L. Zhang, W. Jin, X.Z. Dong, Z.S. Zhao, and X.M. Duan, *Asymmetric fishnet metamaterials with strong optical activity,* Opt. Express **20**, 10776-10787 (2012).

[36] D.M. Bird, and A.R. Preston, *Observation of Berry's geometrical phase in electron diffraction from a screw dislocation*, Phys. Rev. Lett. **61**, 2863-2866 (1988).

[37] A. Mesaros, D. Sadri, and J. Zaanen, *Parallel transport of electrons in graphene parallels gravity*, Phys. Rev. B **82**, 073405 (2010).

[38] M.A.H. Vozmediano, M.I. Katsnelson, and F. Guinea, *Gauge fields in graphene*, Phys. Rep. **496**, 109-148 (2010).